

## CXCL 13 und andere Chemokine

**Ektope Expression von CXCL13 in Gewebe, Blut und Liquor. CXCL13 mittels ELISA. Der vermutete Cutoff für Serum CXCL13 Werte liegt bei 105-115 pg/ml.**

[http://www.arztwww.at/2009\\_10/infekt.html#01](http://www.arztwww.at/2009_10/infekt.html#01)

**Ectopic expression of CXCL13 in tissue, blood, and cerebrospinal fluid. CXCL13 by ELISA. The presumed cutoff values for serum CXCL13 is 105-115 pg / ml.**

### CXCL 13

Ansel KM, Ngo VN, Hyman PL, et al. (2000). "A chemokine-driven positive feedback loop organizes lymphoid follicles". *Nature* 406 (6793): 309–14. [doi:10.1038/35018581](https://doi.org/10.1038/35018581).

Ansel KM, Harris RB, Cyster JG (2002). "CXCL13 is required for B1 cell homing, natural antibody production, and body cavity immunity". *Immunity* 16 (1): 67–76. [doi:10.1016/S1074-7613\(01\)00257-6](https://doi.org/10.1016/S1074-7613(01)00257-6).

Rupprecht TA, Pfister HW, Angele B, (2005) The chemokine CXCL13 (BLC): A putative diagnostic marker for neuroborreliosis. *Neurology* 65(3):448-450.

Narayan K, Dail D, Li L et al. (2005) [The nervous system as ectopic germinal center: CXCL13 and IgG in lyme neuroborreliosis](#). *Ann Neurol.* 57(6), 813-23.

Rupprecht TA, Koedel U, Angele B, et al. (2006) Cytokine CXCL13 - a possible early CSF marker for neuroborreliosis. *Nervenarzt* 77(4), 470-473.

Krumbholz M, Theil D, Cepok S, et al. (2006) Chemokines in multiple sclerosis: CXCL12 and CXCL13 up-regulation is differentially linked to CNS immune cell recruitment. *Brain* 129, 200–211. [Find this article online](#)

Meinl E, Krumbholz M, Hohlfeld R. (2006) lineage cells in the inflammatory CNS environment: Migration, maintenance, local antibody production, and therapeutic modulation. *Ann.Neurol.* 59, 880-892

de Leval L, Rickman DS, Thielen C, et al. (2007). "The gene expression profile of nodal peripheral T-cell lymphoma demonstrates a molecular link between angioimmunoblastic T-cell lymphoma (AITL) and follicular helper T (TFH) cells". *Blood* 109 (11), 4952–63. [doi:10.1182/blood-2006-10-055145](https://doi.org/10.1182/blood-2006-10-055145).

Rupprecht TA, Kirschning CJ, Popp B et al. (2007) [Borrelia garinii induces CXCL13 production in human monocytes through Toll-like receptor 2](#). *Infect Immun* 75(9), 4351-4356.

Ljostad U, Mygland A. (2008) CSF B - lymphocyte chemoattractant (CXCL13) in the early diagnosis of acute Lyme neuroborreliosis. *J Neurol* Epub ahead of print.

King M, Poya H, Rao J, Natarajan Set al. (2009) [CXCL13 expression in Chlamydia trachomatis infection of the female reproductive tract](#). *Drugs Today (Barc).* 45 Suppl B:125-34.

Rupprecht TA, Plate A, Adam M et al. (2009) [The chemokine CXCL13 is a key regulator of B cell recruitment to the cerebrospinal fluid in acute Lyme neuroborreliosis](#). *J Neuroinflammation.* 30;6:42. [Free PMC Article](#)

Sellebjerg F, Bornsen L, Khademi M, et al. (2009) Increased cerebrospinal fluid concentrations of the chemokine CXCL13 in active MS. *Neurology* 73: 2003–2010. [Find this article online](#)

Sellebjerg F, Börnsen L, Khademi M et al. (2009) [Increased cerebrospinal fluid concentrations of the chemokine CXCL13 in active MS](#). *Neurology.* 73(23):2003-10.

- Wutte N (2009) CXCL13 as a diagnostic marker for Lyme Neuroborreliosis. Diploma Thesis. [https://www.google.de/search?q=Diagnostic+value+of+cytokines+and+chemokines+in+Lyme+neuroborreliosis&ie=utf-8&oe=utf-8&rls=org.mozilla:de:official&client=firefox-a&gws\\_rd=cr](https://www.google.de/search?q=Diagnostic+value+of+cytokines+and+chemokines+in+Lyme+neuroborreliosis&ie=utf-8&oe=utf-8&rls=org.mozilla:de:official&client=firefox-a&gws_rd=cr)
- Patadia M, Dixon J, Conley D et al. (2010) [Evaluation of the presence of B-cell attractant chemokines in chronic rhinosinusitis](#). Am J Rhinol Allergy. 24(1):11-6.
- Senel M, Rupprecht TA, Tunami H, Pfister HW, Ludolph AC, Brettschneider J (2010) The chemokine CXCL13 in acute neuroborreliosis. J Neurol Neurosurg Psychiatry 81, 929-933.
- Zhong X, Wang H, Dai Y, Wu A, et al. (2011) [Cerebrospinal fluid levels of CXCL13 are elevated in neuromyelitis optica](#). J Neuroimmunol. 15;240-241:104-8. PMID 22036953.
- EI-Haibi CP, et al. (2011) [CXCL13 mediates prostate cancer cell proliferation through JNK signalling and invasion through ERK activation](#). Cell Prolif, PMID 21645150.
- Del Grosso F, et al. (2011) [Role of CXCL13-CXCR5 crosstalk between malignant neuroblastoma cells and Schwannian stromal cells in neuroblastic tumors](#). Mol Cancer Res, PMID 21642390.
- Ezzat MHM, EL-Gammasy TMA, Shaheen KYA et al. (2011) [Elevated production of serum B-cell-attracting chemokine-1 \(BCA-1/CXCL13\) is correlated with childhood-onset lupus disease activity, severity, and renal involvement](#). Lupus 20(8): 845-854 [Abstract](#)
- Ragheb S, et al. (2011) [Multiple sclerosis: BAFF and CXCL13 in cerebrospinal fluid](#). Mult Scler, PMID 21372118.
- Khademi M, Kockum I, Andersson ML et al. (2011) [Cerebrospinal fluid CXCL13 in multiple sclerosis: a suggestive prognostic marker for the disease course](#). Mult Scler. 17(3):335-43.
- Tumani H, Cadavid D. (2011) Are high CSF levels of CXCL13 helpful for diagnosis of Lyme neuroborreliosis? Neurology 76, 1034-1035 <http://www.neurology.org/content/76/12/1051.abstract>
- van Burgel ND, Bakels F, Kroes ACM et al. (2011) Discriminating Lyme Neuroborreliosis from Other Neuroinflammatory Diseases by Levels of CXCL13 in Cerebrospinal Fluid J. Clin. Microbiol. 49(5), 2027-2030 [Abstract](#)
- EI-Haibi CP, Singh R, Sharma PK et al. (2011) [CXCL13 mediates prostate cancer cell proliferation through JNK signalling and invasion through ERK activation](#). Cell Prolif. 44(4), 11-9.
- Schiffer L, Kielstein JT, Haubitz M et al. (2011) Elevation of serum CXCL13 in SLE as well as in sepsis Lupus 20(5), 507-511 [Abstract](#)
- Rosengren S, Wei N, Kalunian KC et al. (2011) CXCL13: a novel biomarker of B-cell return following rituximab treatment and synovitis in patients with rheumatoid arthritis Rheumatology (Oxford) 50(3), 603-610 [Abstract](#)
- Van Burgel ND, Bakels F, Kroes ACM, van Dam AP (2011) Discriminating Lyme Neuroborreliosis from other Neuroinflammatory diseases by levels of CXCL13 in cerebrospinal fluid. J Clin Microbiol 49, 2027-2030.
- Schmidt C, Plate A, Angele B, Pfister H-W, Wick M, Koede U, Rupprecht TA (2011) A prospective study on the role of CXCL13 in Lyme neuroborreliosis. Neurology 76, 1051-1058.
- Wutte N, Berghold A, Aberer E (2011) [CXCL13 chemokine in pediatric and adult neuroborreliosis](#). Acta Neurol Scand 124(5), 321-8
- Wutte N, Berghold A, Krainberger I, et al. (2011) Serum CXCL13 chemokine is not a marker for active Lyme borreliosis. Acta Derm Venereol 91(6), 724-5. [Full Citation](#)

Tjernberg I, Henningsson AJ, Ernerudh J (2011) [Diagnostic performance of cerebrospinal fluid chemokine CXCL13 and antibodies to the C6-peptide in Lyme neuroborreliosis](#). J Infect 62(2), 149-58

Zajkowska J, Moniuszko-Malinowska A\*, Pancewicz SA, Muszyńska-Mazur A, Kondrusik M, Grygorczuk S, Świerzińska-Pijanowska R, Dunaj J, Czupryna P (2011) Evaluation of CXCL10, CXCL11, CXCL12 and CXCL13 chemokines in serum and cerebrospinal fluid in patients with tick borne encephalitis (TBE). Advances in Medical Sciences 56, 311-317 DOI: 10.2478/v10039-011-0033-z

Bugatti S, Manzo A, Benaglio F. et al. (2012) Serum levels of CXCL13 are associated with ultrasonographic synovitis and predict power Doppler persistence in early rheumatoid arthritis treated with non-biological disease-modifying anti-rheumatic drugs. Arthritis Research & Therapy 14:R34 doi:10.1186/ar3742 <http://arthritis-research.com/content/14/1/R34/abstract>

Baay-Guzman GJ, Huerta-Yepe S, Vega MI, et al. (2012) Role of CXCL13 in Asthma: Novel Therapeutic Target. Chest 141, 4 886-894

Kowarik MC, Cepok S, Sellner J et al. (2012) CXCL13 is the major determinant for B cell recruitment to the CSF during neuroinflammation Journal of Neuroinflammation 9,93 doi:10.1186/1742-2094-9-93

Cerar T et al. (2013) Diagnostic value of cytokines and chemokines in Lyme neuroborreliosis Clin Vaccine Immunol. [http://scibite.com/site/library/2013\\_8/1/0/23945160.html](http://scibite.com/site/library/2013_8/1/0/23945160.html)

**Unbound MEDLINE results for: CXCL13 AND human** [Refine this search](#)

321 journal articles in the PubMed database

[See all \(81\) citations in PubMed](#)

## Andere Chemokine

Geeta Ramesh G (2014) Novel Therapeutic Targets in Neuroinflammation and Neuropathic Pain. Inflamm Cell Signal 1, 139-147. doi: 10.14800/ics.111.

<http://www.smartscitech.com/index.php/ICS/article/view/111>

**„Together, CCL2/CCR2 and MAPK signaling networks represent novel therapeutic targets for the development of therapeutic strategies aimed at limiting neuroinflammation and neuropathic pain associated with neurological disease states.“**

Bernt - Dieter Huismans. Letzte Revision Juli 2014 [www.Huismans.de.vu](http://www.Huismans.de.vu), [www.Huismans.de.nu](http://www.Huismans.de.nu)

