

Lichen sclerosus, Lichenoid dermatitis

[Aberer E](#), [Kollegger H](#), et al. (1988) **Neuroborreliosis in morphea and lichen sclerosus et atrophicus**. *JAAD* 19(5), 1, 820-825 [http://www.jaad.org/article/S0190-9622\(88\)70239-X/abstract](http://www.jaad.org/article/S0190-9622(88)70239-X/abstract)

Schempp C, Bocklage H, Lange R, Kölmel HW, Orfanos CE, Gollnick H. (1993) **Further evidence for *Borrelia burgdorferi* infection in morphea and lichen sclerosus et atrophicus confirmed by DNA amplification**. *J Invest Dermatol.* 100(5), 717-20.

Trevisan G, Rees DH, Stinco G. (1994) ***Borrelia burgdorferi* and localized scleroderma**. *Clin Dermatol.* 12(3), 475-9.

Powell JJ, Wojnarowska F (1999) **Lichen sclerosus**. *Lancet.* 353, S. 1777–1783.

Schempp C, Bocklage H, Lange R et al. (1993) **Further evidence for *Borrelia burgdorferi* infection in morphea and lichen sclerosus et atrophicus confirmed by DNA amplification**. *The Journal of investigative dermatology*. Band 100, Nummer 5, S. 717–720, [ISSN 0022-202X](#). [PMID 8491994](#).

Colomé-Grimmer MI, Payne DA, Tying SK, Sánchez RL. (1997) ***Borrelia burgdorferi* DNA and *Borrelia hermsii* DNA are not associated with morphea or lichen sclerosus et atrophicus in the southwestern United States**. *Acta Derm Venereol.* 77(4), 299-304.

Fujiwara H, Fujiwara K, Hashimoto K, et al. (1997) **Detection of *Borrelia burgdorferi* DNA (*B. garinii* or *B. afzelii*) in morphoea and lichen sclerosus et atrophicus tissues of German and Japanese but not of US patients**. *Arch Dermatol* 133, 41–4.

Svecova D, Buchvald J. (2000) ***Borrelia burgdorferi* antibodies in scleroderma circumscripta, lichen sclerosus et atrophicus, erythema nodosum, granuloma annulare, erythema annulare and chronic urticaria**. *Int J Dermatol.* 39(4) 278-83.

Ozkan S, Atabey N, Fetil E, Erkizan V, Günes AT. (2000) **Evidence for *Borrelia burgdorferi* in morphea and lichen sclerosus**. *Int J Dermatol.* 39(4), 278-83.

[Breier F](#), [Khanakah G](#), [Stanek G](#), [Kunz G](#), [Aberer E](#), [Schmidt B](#), [Tappeiner G](#). (2001) **Isolation and polymerase chain reaction typing of *Borrelia afzelii* from a skin lesion in a seronegative patient with generalized ulcerating bullous lichen sclerosus et atrophicus**. *Br J Dermatol.* 144(2), 387-92. <http://www.ncbi.nlm.nih.gov/pubmed/11251580>

Kaya G, Berset M, Prins C, Chavaz P, Saurat JH. (2001) **Chronic borreliosis presenting with morphea- and lichen sclerosus et atrophicus-like cutaneous lesions. A case report**. *Dermatology.* 202(4), 373-5

Shelley WB, Shelley ED, Amurao CV. (2006) **Treatment of lichen sclerosus with antibiotics**. *Int J Dermatol.* 45(9), 1104-6.

Eisendle K, Grabner T, Kutzner H et al. (2008) **Possible Role of *Borrelia burgdorferi* Sensu Lato Infection in Lichen Sclerosus**. *Archives of Dermatology.* 144, S. 591–598, [doi:10.1001/archderm.144.5.591](https://doi.org/10.1001/archderm.144.5.591).

[Cesinaro AM](#), [Sighinolfi P](#), [Greco A](#) et al. (2009) **Annular lichenoid dermatitis of youth ... and beyond: a series of 6 cases**. *Am J Dermatopathol.* 31(3), 263-7. doi: 10.1097/DAD.0b013e31819af679. <http://www.ncbi.nlm.nih.gov/pubmed/19384067>

Edmonds E, Mavin S, Francis N, Ho-Yen D, Bunker C. (2009) ***Borrelia burgdorferi* is not associated with genital lichen sclerosus in men**. *Br J Dermatol.* 160(2), 459-60

Zollinger T, Mertz KD, Schmid M et al. (2010) ***Borrelia* in granuloma annulare, morphea and lichen sclerosus: a PCR-based study and review of the literature**. *Journal of Cutaneous Pathology.* 37, 571–577, [doi:10.1111/j.1600-0560.2009.01493.x](https://doi.org/10.1111/j.1600-0560.2009.01493.x).

Susana Aid S, Lattario FR et al. (2010) **Epstein-Barr Virus and Human Papillomavirus Infection in Vulvar Lichen Sclerosus**. Journal of Lower Genital Tract Disease. 14, 319–322, doi:10.1097/LGT.0b013e3181d734f1.

Belz Doreen (2013) **Expression von regulatorischen T-Zellen beim genitalen Lichen sclerosus**. Inaugural Dissertation , Ruhr Universität Bochum. <http://www-brs.ub.ruhr-uni-bochum.de/netahtml/HSS/Diss/BelzDoreen/diss.pdf>

Kirtschig G, Cooper S, Aberer W et al. (2015) **Evidence based (S3) guideline on (anogenital) Lichen sclerosus**. J Eur Acad Dermatol Venerol 29, e1-43 doi: 10.1111/jdv.13136. <https://www.ncbi.nlm.nih.gov/pubmed/26202852>
http://www.sggg.ch/fileadmin/user_upload/Dokumente/3_Fachinformationen/2_Guidelines/De/Guideline_on_Lichen_sclerosus.pdf

Kirtschig G (2016) **Lichen sclerosus—presentation, diagnosis and management**. Dtsch Arztebl Int 2016; 113: 337–43. DOI: 10.3238/arztebl.2016.0337 <http://www.aerzteblatt.de/archiv/178786>

Wilk M, Zelger B, Emberger M et al. (2016) **Annular Lichenoid Dermatitis (of Youth) Immunohistochemical and Serological Evidence of Another Clinical Presentation of Borrelia Infection in Patients of West Austria**. Am J of Dermatology. doi: 10.1097/DAD.0000000000000621 http://journals.lww.com/amjdermatopathology/Abstract/publishahead/Annular_Lichenoid_Dermatitis_of_Youth_.98764.aspx
„Lichen sclerosus et atrophicus and morphea have previously been reported to be possibly related to Borrelia infection. We postulate that a similar relationship to Borrelia infection may be true for ALDY implying that ALDY may be an early superficial stage of morphea.“

[Bernt - Dieter Huismans](#), Start 2016. Letzte Revision Januar 2017 www.Huismans.click
Back to top: http://www.erlebnishaft.de/lichen_sclerosus.pdf

