

LTT (Lymphozyten-Proliferationstest, „Lymphozyten-Transformationstest“)

Interferon Gamma Test, ELISpot (T-Cell Spot), dem Lymphozytentransformationstest bei Borreliose und bei anderen Infektionskrankheiten

Im **Lymphozyten-Proliferationstest**, auch **Lymphozyten-Transformationstest** genannt (**LTT**), wird die **Proliferationskapazität von Lymphozyten** dargestellt. Der Test dient der Bewertung des Immunstatus oder der Diagnose einer speziellen Sensibilisierung.

In the **Lymphocyte proliferation assay**, also known as **lymphocyte transformation test (LTT)**, the **proliferation capacity of lymphocytes** is shown. The test is used to evaluate the immune status or the diagnosis of specific sensitization of an object.

Im **Interferon Gamma Test**, dem **ELISPOT (Enzyme-Linked ImmunoSpot)**, auch T-Cellspot genannt, wird durch die **Bestimmung der Zytokin – Produktion die Quantität und die Qualität einer T – Zell - Immunantwort zeitnah** dokumentiert.

In the **Interferone Gamma Test**, **ELISPOT (Enzyme-Linked ImmunoSpot)**, also called T-Cell spot, represented by the determination of cytokine – Production, the **quantity and quality of T - cell - immune response in a timely manner** is documented.

Lim LC, England DM, DuChateau BK, Glowacki NJ, Schell RF (1995) **Borrelia burgdorferi-specific T lymphocytes** induce severe destructive Lyme arthritis. Infect. Immun. 63, 1400-1408.

Dwyer JM et al. (1979) Behavior of human immunoregulatory cells in culture. I Variables requiring consideration for clinical studies. Clin Exp Immunol 38, 499-513

Johnson C, Dwyer JM (1981) Quantitation of spontaneous suppressor cell activity in human cord blood lymphocytes and their behavior in culture (abstract). Fed Proc 40, 1075

Czerkinsky C, Nilsson L, Nygren H, Ouchterlony O, Tarkowski A (1983) **A solid-phase enzyme-linked immunospot (ELISPOT) assay for enumeration of specific antibody-secreting cells.** J Immunol Methods 65 (1–2), 109–121. doi:10.1016/0022-1759(83)90308-3. PMID 6361139 <http://www.ncbi.nlm.nih.gov/pubmed/6361139>

Sigal LH et al (1984) Cellular immune findings **Lyme disease**. Yale J Biol Med 57, 595-8

Sigal LH et al (1986) Proliferative responses of mononuclear cells in **Lyme disease**. Reactivity to Borrelia burgdorferi antigens is greater in joint fluid than in blood. Arthritis Rheum 29, 761-9

Dattwyler RJ et al (1986) Cellular immune response in **Lyme disease**: the response to mitogens, I live Borrelia burgdorferi, NK cell function and lymphocyte subsets. Zentralbl Bakteriell Mikrobiol Hyg 263(1-2), 151-9

Dattwyler RJ et al (1988) Seronegative **Lyme disease**. Dissociation of specific T- and B-lymphocyte responses to Borrelia burgdorferi. N Engl J Med 319, 1441-6

Dattwyler RJ, Volkman, DJ, Halperin JJ, Luft BJ, Thomas J, Golightly MG (1988) Specific immune responses in **Lyme borreliosis**. Characterization of T cell and B cell responses to Borrelia burgdorferi. Ann. N. Y Acad. Sci. 539, 93-102

Mosmann TR et al (1989) TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol.* 7, 145-73

[Dressler F](#), [Yoshinari NH](#), [Steere AC](#). (1991) The T-cell proliferative assay in the diagnosis of **Lyme disease**. *Ann Intern Med.* 115(7), 533-9 <http://www.ncbi.nlm.nih.gov/pubmed/1883122>

Krause et al (1991) T cell proliferation induced by *Borrelia burgdorferi* in patients with **Lyme borreliosis**. Autologous serum required for optimum stimulation. *Arthritis Rheum* 34 393-402

Zoschke DC et al (1991) Lymphoproliferative responses to *Borrelia burgdorferi* in **Lyme disease**. *Ann Intern med* 17, 151-8

Schempp et al (1993) Comparison of *Borrelia burgdorferi* ultrasonicate and whole **B.burgdorferi** cells as a stimulus for T-cell proliferation and GM-CSF secretion in vitro. *Zentralbl Bakteriologie* 279, 417-25

Horowitz HH et al (1994) **Sustained cellular immune responses to *Borrelia burgdorferi*: lack of correlation with clinical presentation and serology**. *Clin Diagn Lab Immunol* 1, 373-8

Roessner, K., Russell, J. Q., Cooper, S. M., Budd, R. C., Fikrig, E., & Flavell, R. A. (1994) Prominent T lymphocyte response to *Borrelia burgdorferi* from peripheral blood of unexposed donors. *European journal of immunology* 24(2), 320-324

Forsberg P, Ernerudh J, Ekerfelt C et al (1995) The outer surface proteins of **Lyme disease** *Borrelia* spirochetes stimulate T cells to secrete interferon gamma (IFN-Gamma): diagnostic and pathogenic implications. *Clinical and Experimental Immunology* 101(3), 453-460

Breier, F. et al. (1995) Lymphocyte proliferation test in cutaneous manifestations of **Lyme Borreliosis**. *Wien Med Wochenschr.* 145 (7-8): 170-3.

Buechner SA et al (1995) Lymphoproliferative responses to ***Borrelia burgdorferi*** in patients with erythema migrans, acrodermatitis chronica atrophicans, lymphadenitis benigna cutis and morphea. *Arch Dermatol* 131, 673-7

Breier F et al (1996) Lymphoproliferative responses to ***Borrelia burgdorferi*** in circumscribed scleroderma. *Brit J Dermatol* 134, 285-91

Huppertz et al (1996) Lymphoproliferative responses to ***Borrelia burgdorferi*** in the diagnosis of Lyme arthritis in children and adolescents. *Eur J Pediatr* 155, 297-302

Rutkowski S et al (1997) Lymphocyte proliferation assay in response to ***Borrelia burgdorferi*** in patients with Lyme arthritis: analysis of lymphocyte subsets. *Rheumatol Int* 17, 151-8

Ekerfeldt C, Ernerudh J, Bunikis J et al (1997) Compartmentalization of antigen specific cytokine response to the central nervous system in CNS **borreliosis**: secretion of IFN-Gamma predominates over IL-4 secretion in response to outer surface proteins of lyme disease ***Borrelia spirochetes***. *Journal of Neuroimmunology* 79(2), 155-162

Fendler, C. et al (1998) Longitudinal Investigation of Bacterium-Specific Synovial Lymphocyte Proliferation In Reactive Arthritis and **Lyme Arthritis**. *Br J Rheumatol* 37, 784-788.

Ekerfelt C, Forsberg P, Senvik M et al. (1999) Asymptomatic *Borrelia*-seropositive individuals display the same incidence of ***Borrelia*-specific interferon-gamma (IFN-g)-secreting cells** in blood as patients with clinical *Borrelia* infection. *Clin Exp Immunol.* 115, 498-502.

Baehr von V et al (2001) Improving the in vitro antigen specific T- cell proliferation assay: the use of interferon alpha to elicit antigen specific stimulation and decrease bystander proliferation. *J. Immunol. Methods* 251, 63- 71

Pohl-Koppe A et al (2001) Charakterization of the cellular and humoral immune response to outer surface protein C and outer surface protein 17 in children with early disseminated **Lyme borreliosis**. *Med Microbiol Immunol* 189, 193-200

Lavani A, Pathan AA, Durkan H, Wilkinson KA, Whelan A, Deeks JJ, Reece WH, Latif M, Pasvol, G, Hill AV (2001) Enhanced contact tracing and spatial tracking of **Mycobacterium tuberculosis infection** by enumeration of antigen-specific T cells. *Lancet* 357, 2017-2021.

Bauer, Y. et al (2001) Prominent T cell response to a selectively in vivo expressed **Borrelia burgdorferi** outer surface protein (pG) in patients with Lyme disease. *Eur. J. Immunol.* 31, 767-776

Vaz A et al (2001) Cellular and humoral immune responses to **Borrelia burgdorferi** antigens in patients with culture-positive early Lyme disease. *Infect Immun* 69, 7437-44

Bundesgesundheitsbl - Gesundheitsforsch - Gesundheitsschutz (2002) 45, 745–749 DOI 10.1007/s00103-002-0469-1 **Diagnostische Relevanz des Lymphozytentransformationstestes in der Umweltmedizin Mitteilung der Kommission „Methoden und Qualitätssicherung in der Umweltmedizin“ am Robert Koch-Institut (RKI)**
http://edoc.rki.de/documents/rki_ab/re67fIHRghoUo/PDF/26C0oqlxZYpo.pdf

Gerritzen A et al (2004) Lymphocyte Transformation Test (LTT) as a helpful diagnostic tool to detect active **Lyme Borreliosis** *Int. J. Medical Microbiology* 294S1, 196

Lavani A (2004) Counting antigen-specific T cells: A new approach for **monitoring response to tuberculosis treatment?** *Clin. Infect. Dis.* 38, 757-759

Wilske B und Fingerle V (2005) **Lyme-Borreliose** *Diagnostik. Mikrobiologie* 15.Jg.

Kalyunzhny AE (2005) Handbook of **ELISPOT** Methods and Protocols.
http://books.google.de/books?id=IAMBUFWHgMwC&pg=PA27&lpg=PA27&dq=A+solid-phase+enzyme-linked+immunospot+%28ELISPOT%29+assay+for+enumeration+of+specific+antibody-secreting+cells&source=bl&ots=OnzIBAY7ew&sig=mm-BIQPfm8f-nUhHQjbYSsvFVI&hl=de&sa=X&ei=p5W2U4L9GamJ4qS_soCgBw&ved=0CEAQ6AEwAw#v=onepage&q=A%20solid-phase%20enzyme-linked%20immunospot%20%28ELISPOT%29%20assay%20for%20enumeration%20of%20specific%20antibody-secreting%20cells&f=false

Ibd. Chemistry and Biology of the ELISPOT Assay

http://books.google.de/books?id=IAMBUFWHgMwC&pg=PA27&lpg=PA27&dq=A+solid-phase+enzyme-linked+immunospot+%28ELISPOT%29+assay+for+enumeration+of+specific+antibody-secreting+cells&source=bl&ots=OnzIBAY7ew&sig=mm-BIQPfm8f-nUhHQjbYSsvFVI&hl=de&sa=X&ei=p5W2U4L9GamJ4qS_soCgBw&ved=0CEAQ6AEwAw#v=onepage&q=A%20solid-phase%20enzyme-linked%20immunospot%20%28ELISPOT%29%20assay%20for%20enumeration%20of%20specific%20antibody-secreting%20cells&f=false

Jarefors S et al (2006) Reduced number of interleukin-12 secreting cells in patients with Lyme borreliosis previously exposed to *Anaplasma phagocytophilum*. *Clin Exp Immunol.* 143, 322-8

Valentine-Thon E et al (2007) A novel lymphocyte transformation test for **Lyme borreliosis**. *Diagn Microbiol Infect Dis* 57, 27-34

Von Baehr V et al (2007) Untersuchungen zur diagnostischen Wertigkeit des Lymphozytentransformationstestes bei Patienten mit **Borreliose**. *J Lab Med* 31(3), 149-158
<http://www.degruyter.com/dg/viewarticle/j%24002flabm.2007.31.issue-3%24002film.2007.023%24002film.2007.023.xml>

„Qualitätssicherung beim Lymphozytentransformationstest“ – Addendum zum LTT-Papier der RKI-Kommission „Methoden und Qualitätssicherung in der Umweltmedizin“ Mitteilung der Kommission „Methoden und Qualitätssicherung in der Umweltmedizin“ Bundesgesundheitsbl - Gesundheitsforsch - Gesundheitsschutz 2008 · 51:1070–1076 DOI 10.1007/s00103-008-0641-3 Online publiziert: 7. September 2008 © Springer Medizin Verlag 2008
http://edoc.rki.de/documents/rki_ab/re67fIHRghoUo/PDF/26C0oxyqlxZYff.pdf

Gerritzen, A. (2008) Verbesserte **Borrelien-Serologie** durch Einsatz von rekombinanten Antigenen mit der luminex-Technologie. *Borreliose Wissen* Nr. 17, 14-16

Valentine-Thon E, Ilsemann K, Sandkamp M (2008) **Neuartiger Lymphozyten-Transformations-Test (LTT-MELISA®) zum Nachweis einer Lyme-Borreliose / A novel lymphocyte transformation test (LTT-MELISA®) for Lyme borreliosis**. LaboratoriumsMedizin. Band 32, Heft 1, Seiten 26–34, ISSN (Online) 1439-0477, ISSN (Print) 0342-3026, DOI: [10.1515/JLM.2008.007](https://doi.org/10.1515/JLM.2008.007), January 2008 <http://www.degruyter.com/view/j/jlm.2008.32.issue-1/jlm.2008.007/jlm.2008.007.xml?format=INT>

Zhang W, Tary-Lehmann M (2008) Objective, User-Independent **ELISPOT** Data Analysis Based on Scientifically Validated Principles. In Validation of **Cell-Based Assays** in the GLP Setting: A Practical Guide, 1st ed.; Uma Prabhakar, Marian Kelley, Eds.; John Wiley & Sons, Ltd.: West Sussex, UK, 126-147.

Tary-Lehmann M, Hamm CD, Lehmann PV. (2008) Validating reference samples for comparison in a regulated **ELISPOT** assay. In: Uma Prabhakar and Marian Kelley Eds. Validation of Cell-Based Assays in the GLP Setting: A Practical Guide. 1st Ed. West Sussex, England: John Wiley & Sons. Ltd; 127-146

Zhang W, Caspell R, Karulin AY, Ahmad M, Haicheur N, Abdelsalam A, Johannesen K, Vignard V, Dudzik P, Georgakopoulou K, et al. (2009) **ELISPOT** assays provide reproducible results among different laboratories for T-cell immune monitoring--even in hands of **ELISPOT**-inexperienced investigators. 1 Immunotoxicol. 6, 227-234

Berghoff W (2009) **Lymphozytentransformationstest (LTT) bei der Lyme-Borreliose Literaturübersicht**. http://www.praxis-berghoff.de/dokumente/LTT_bei_der_Lyme-Borreliose.pdf

CDC (2010) Updated Guidelines for Using Interferon Gamma Release Assays to Detect Mycobacterium tuberculosis Infection --- United States. Recommendations and Reports 59(RR05), 1-25 (ELISPOT) http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5905a1.htm?s_cid=rr5905a1_e

CDC (2011) TB Elimination. Interferon-Gamma Release Assays (IGRAs) – Blood Tests for TB Infection <http://www.cdc.gov/tb/> <http://www.cdc.gov/tb/publications/factsheets/testing/IGRA.pdf>

T-Cell-Spot/IGRA has been approved by the FDA in May 2011 for M. tuberculosis; TB EliSpot (TSpot TB) by Oxford Immunotech. www.fda.com PMA (premarket approval) nr. P070006

Nordberg et al. (2012) Can **ELISPOT** be applied to a clinical setting as a diagnostic utility for Neuroborreliosis?, Cells I, 153-167

Kuerten, S., Batoulis, H., Recks, M.S., et al. (2012) Resting of Cryopreserved PBMC Does Not Generally Benefit the Performance of Antigen-Specific T Cell **ELISPOT** Assays. Cells 1(3), 409-427

Lehman PV et al. (2012) Unique Strengths of **ELISPOT** for T Cell Diagnostics in: Kalyuzhny AE. Handbook of ELISPOT: Methods and Protocols, Methods in Molecular Biology, Vol. 792. 2nd Ed: Springer 3-23

von Baehr V et al. (2012) **The Lymphocyte Transformation Test for Borrelia Detects Active Lyme Borreliosis and Verifies Effective Antibiotic Treatment**, The Open Neurology Journal 6, 104-112
„**Borrelia-specific antibodies are not detectable until several weeks after infection and even if they are present, they are no proof of an active infection. Since the sensitivity of culture and PCR for the diagnosis or exclusion of borreliosis is too low, a method is required that detects an active Borrelia infection as early as possible. For this purpose, a lymphocyte transformation test (LTT) using lysate antigens of Borrelia burgdorferi sensu stricto, Borrelia afzelii and Borrelia garinii and recombinant OspC was developed and validated through investigations of seronegative and seropositive healthy individuals as well as of seropositive patients with clinically manifested borreliosis. The sensitivity of the LTT in clinical borreliosis before antibiotic treatment was determined as 89,4% while the specificity was 98,7%. In 1480 patients with clinically suspected borreliosis, results from serology and LTT were comparable in 79.8% of cases. 18% were serologically positive and LTT-negative. These were mainly patients with borreliosis after antibiotic therapy. 2.2% showed a negative serology and a positive LTT result. Half of them had an early erythema migrans. Following antibiotic treatment, the LTT became negative or borderline in patients with early manifestations of borreliosis, whereas in patients with late symptoms, it showed a regression while still remaining positive. Therefore, we propose the follow-up monitoring of disseminated Borrelia infections as the main indication for the Borrelia-LTT.**“

Berghoff W (2012) **Handbuch Neurologie 2012, Neuroborreliose. Kritische Stellungnahme „Fazit für die Praxis: Der LTT wird aufgrund von Spezifitätsproblemen in der Diagnostik der Lyme-Borreliose/Neuroborreliose nicht empfohlen. Es gibt bislang keine wissenschaftlich gesicherten Daten, die einen Nutzen des LTT/Elispot belegen und den Einsatz in der Diagnostik der Borreliose rechtfertigen. Der Test ist insbesondere auch nicht geeignet, die Krankheitsaktivität einer Borrelieninfektion anzuzeigen; aus dem Testergebnis darf keine Therapieentscheidung abgeleitet werden. Zusammengefasst ist der Borrelien-LTT als Diagnoseinstrument und als Verlaufsparemeter in der Borreliendiagnostik ungeeignet.“**

Skogman BH, Hellberg S, Ekerfeld Ch et al. (2012) Research Article. Adaptive and Innate Immune Responsiveness to **Borrelia burgdorferi** sensu lato in Exposed Asymptomatic Children and Children with Previous Clinical Lyme Borreliosis. Clinical and Developmental Immunology Volume 2012, Article ID 294587,10 pages. doi:10.1155/2012/294587 <http://www.hindawi.com/journals/cdi/2012/294587/>

[Lehmann PV](#), [Zhang W](#). (2012) Unique strengths of **ELISPOT** for T cell diagnostics. Methods Mol Biol. 792, 3-23. doi: 10.1007/978-1-61779-325-7_1. <http://www.ncbi.nlm.nih.gov/pubmed/21956497>

Nordberg M, Forsberg P, Nyman D, Skogman BH, Nyberg C, Ernerudh J, Eliasson I, Ekerfeldt C (2012) Can **ELISPOT** be applied to a clinical setting as a diagnostic utility for Neuroborreliosis? Cells 1, 153-167.

Hagen J, Houchins JP, Kalyuzhny AE (2012) Combining **ELISPOT** and ELISA to measure amounts of cytokines secreted by a single cell. In Handbook of **ELISPOT**: Methods and Protocols, Methods in Molecular Biology, 2nd ed.; Kalyuzhny, A., Ed.; Springer Science+Business Media, LLC: New York, NY, USA, 792, 115-122

Lehmann PV, Zhang W (2012) Unique Strengths of **ELISPOT** for T Cell Diagnostics. In Handbook of **ELISPOT**: Methods and Protocols, Methods in Molecular Biology, 2nd ed.; Kalyuzhny, A., Ed.; Springer Science+Business Media, LLC: New York, NY, USA, 792, 3-23

Chenggang J, Fall DJ, Roen D, Kellermann G (2012) **iSpot Lyme™**: A New Generation of **Lyme Disease Testing**. Neuroscience <https://www.neurorelief.com/index.php?p=cms&cid=486&pid=149>

Chenggang J, Roen DR, Lehmann PV, Kellermann GH (2013) An Enhanced **ELISPOT** Assay for Sensitive Detection of Antigen-Specific T Cell Responses to **Borrelia burgdorferi**. Cells 2, 607-620

Goldman E. (2013) New T-Cell Test a “Game-Changer” for Lyme. Holistic Primary Care. 14(3). <http://holisticprimarycare.net/topics/topics-h-n/infectious-disease/1512-new-t-cell-test-a-game-changer-for-lyme.html>

Bestard O, Crespo E, Stein M et al. (2013) Cross-Validation of **IFN-γ Elispot** Assay for Measuring Alloreactive Memory/Effector T Cell Responses in Renal Transplant Recipients, American J Transplantation XX, 1-11 1880-90. doi: 10.1111/ajt.12285. <http://www.ncbi.nlm.nih.gov/pubmed/23763435>

Bittel P, Mayor D, Iseli P et al. (2014) IGRA-positive patients and interferon-gamma/interleukin-2 signatures: Can the Fluorospot assay provide further information? Infection, Institute for Infectious Diseases, University of Bern. [Infection](#) 42(3), 539-43. doi: 10.1007/s15010-014-0588-2. <http://www.ncbi.nlm.nih.gov/pubmed/24477887>

[Dessau RB](#), [Fingerle V](#), [Gray J](#), [Hunfeld KP](#), [Jaulhac B](#), [Kahl O](#), [Kristoferitsch W](#), [Stanek G](#), [Strle F](#) (2014) **The lymphocyte transformation test for the diagnosis of Lyme borreliosis has currently not been shown to be clinically useful.** Clinical Microbiology and Infection. European Society of Clinical Microbiology and Infectious Diseases. The authors of this publication are members of the ESCMID study group for Lyme borreliosis: www.escmid.org/ESGBOR [Clin Microbiol Infect](#). doi: 10.1111/1469-0691.12583. [Epub ahead of print] <http://www.ncbi.nlm.nih.gov/pubmed/24520912>
„ABSTRACT This letter is a comment on a study using lymphocyte transformation test (LTT) for diagnosis of active Lyme borreliosis caused by Borrelia burgdorferi sensu lato [1]. This LTT study reports the findings derived from a validation panel containing 120 blood donors seronegative for Borrelia, 40 seronegative patients with autoimmune diseases, 48 healthy seropositive controls and 94 seropositive patients with clinical signs of Lyme borreliosis. Furthermore, 1480 samples were investigated with both serology (Borrelia IgG and IgM ELISA and Western blot, Mikrogen, Munich Germany) and LTT.“
„The development of a biomarker for active infection with B. burgdorferi sensu lato would be of clinical value, as antibody detection cannot currently distinguish active infection from immunological memory

resulting from past or asymptomatic infection. However, T-cell recognition may be inherently indiscriminate, and problems with specificity may therefore be hard to avoid. A recent Swedish study [here no citation] “did not find the ELISPOT technique to be useful for supplementary clinical diagnosis, with a specificity of just 82%.... In conclusion, the clinical value of the LTT for the diagnosis of active Lyme borreliosis was not supported the von Baehr et al. Study [1]“

von Baehr V (2014) **The lymphocyte transformation test for the diagnosis of Lyme borreliosis could fill a gap in the difficult diagnostics of borreliosis.** Article first published online: <http://onlinelibrary.wiley.com/doi/10.1111/1469-0691.12755/abstract>
DOI: 10.1111/1469-0691.12755

Berghoff W (2014) Lymphozytentransformationstest (LTT) bei der Lyme-Borreliose. http://www.praxis-berghoff.de/dokumente/berghoff150714/Kapitel_22-6_Lymphozytentransformationstest_LTT.pdf

Berghoff W (2014) Negativbeurteilung des LTT (Auwaerter et al, 2011). http://www.praxis-berghoff.de/dokumente/berghoff150714/Kapitel_22-7_Negativbeurteilung_des_LTT_Auwaerter.pdf

Berghoff W (2014) Korrelation LTT / Serologie. http://www.praxis-berghoff.de/dokumente/berghoff150714/Kapitel_22-8_Korrelation_LTT_Serologie.pdf

von Baehr V (2014) **The lymphocyte transformation test for the diagnosis of Lyme borreliosis.** *Clin Microbiol Infect.* pii: S1198-743X(14)00055-X. doi: 10.1016/j.cmi.2014.10.001. [Epub ahead of print] <http://www.ncbi.nlm.nih.gov/pubmed/25658576>

Puri BK, Segal DR, Monroe JA (2014) **Diagnostic use of the lymphocyte transformation test-memory lymphocyte immunostimulation assay in confirming active Lyme borreliosis in clinically and serologically ambiguous cases.** *Int J Clin Exp Med.* 7(12), 5890-2. eCollection 2014. <http://www.ncbi.nlm.nih.gov/pubmed/25664127>

« It is concluded that, for putative European-acquired Lyme borreliosis infections, it would be sensible to carry out both the LTT-MELISA and Western blot assay. »

BCA-clinic (2015) **EliSpot: Die nächste Generation.** www.infectolab.de
http://www.infectolab.de/wp-content/uploads/2015/05/LymeSpot_PDF_german_print.pdf

Bacher P, Steinbach A, Kniemeyer O et al. (2015) **Fungus-specific CD4(+) T cells for rapid identification of invasive pulmonary mold infection.** *Am J Respir Crit Care Med.*, 191(3), 348-52
doi: 10.1164/rccm.201407-1235LE <http://www.ncbi.nlm.nih.gov/pubmed/25635493>

- ➔ **Elispot Animation** <https://www.mabtech.com/knowledge-center/animations/elispot-animation>
- ➔ **Cytokine Elispot Animation** <http://www.immunospot.eu/elispot-animation.html>
- ➔ **AID iSpot** <http://elispot.com/>
- ➔ **BCA Lyme Spot** (2015) <http://borreliose-centrum-augsburg.de/lymespot/>
<http://borreliose-centrum-augsburg.de/en/lymespot-2/>
<http://borreliose-centrum-augsburg.de/wp-content/uploads/2015/03/LymeSpot-dt.pdf>
- ➔ <http://borreliose-centrum-augsburg.de/wp-content/uploads/2015/03/Info-ohne-Preis-dt-%C3%9C-AID.pdf>

Bernt - Dieter Huismans Letzte Revision November 2016 www.Huismans.click
Back to top: <http://www.erlebnishaft.de/ltt.pdf>

